

CWR FOLIAR NUTRIENT 0-8-0 provides highly available plant food for foliar feeding. It is compatible with organic pesticides for combination spraying.

GUARANTEED ANALYSIS

0-8-0

Available Phosphoric Acid (P ₂ O ₅)	8.0%
Sulfur (S)	2.2%
Iron (Fe)	3.0%
Zinc (Zn)	1.0%

Phosphate derived from phosphoric acid; sulfur, iron and zinc derived from metallic sulfates.

WARNING: This product contains chemicals known to the State of California to cause cancer and birth defects or other reproductive harm.

CAUTION: Causes eye irritation. In case of contact with the eyes, immediately flush with water for at least 15 minutes. If irritation persists, get medical attention. Do not swallow. Do not repack into unlabeled containers, or those normally used for food or drink.


READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS.

Do not store below 32°F.

STATEMENT OF WARRANTY

NOTICE: Seller's guarantee shall be limited to the terms of the label. Subject thereto, the buyer assumes any risk to persons or property arising from the use or handling and accepts the product on these conditions.

FOR AGRICULTURAL SPRAYS



**KEEP OUT OF REACH OF CHILDREN
CAUTION**

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.regulatory-info-monterey.com>

NET CONTENTS: 1 GALLON / 3.8 LITERS

Manufactured by:



CREATIVE MARKETING & RESEARCH, INC.
P.O. Box 35000 • Fresno, CA 93745-5000
(559) 499-2100

* Registered TM of CREATIVE MARKETING & RESEARCH, INC.

DIRECTIONS FOR USE

For row or field crops, mix **CWR FOLIAR NUTRIENT 0-8-0** in at least 20 gallons water per acre for ground sprayers or 10 gallons for aircraft. Apply to fruit trees or ornamentals in at least 100 gallons water per acre with conventional sprayers, 45-100 gallons with concentrate sprayers, or 10 gallons for aircraft. On waxy foliage or fruit, add sufficient spreader for smooth wetting to avoid spotting.

Use quantities indicated in the table below or follow recommendations of local agricultural authorities. Add spreader as needed for smooth wetting.

USE PRECAUTIONS: Follow recommendation of local agricultural authorities for proper timing. **CWR FOLIAR NUTRIENT 0-8-0** is primarily intended for post-bloom use during fruit and seed formation. Make test applications on a few plants if using combinations not known to be safe. Use only in equipment free of herbicide contamination. To avoid phytotoxicity, do not apply with copper or lime-sulfur fungicides, or during bloom.

APPLES, APRICOTS, PEARS, PLUMS, PRUNES, NECTARINES, PEACHES, CHERRIES, ALMONDS, PECANS, WALNUTS, OLIVES, FIGS and AVOCADOS: Apply 2 quarts/acre at 2 to 3 week intervals after bloom. On smooth skinned fruits, add spreader to insure "drop-free" wetting.

ALFALFA, CLOVER: Apply 1 quart/acre as first spring growth starts and repeat after each cutting.

BEANS, BARLEY, CORN, RICE, MILO, WHEAT, OATS, SAFFLOWER and SUGAR BEETS: Apply 1 to 2 quarts/acre in post-bloom applications.

CITRUS: Apply 2 quarts/acre at 2 to 3 week intervals as needed.

ARTICHOKES, CELERY, CARROTS, COLE CROPS, SPINACH and SWEET CORN: Apply 2 quarts/acre as needed.

COTTON: Apply 1 to 2 quarts/acre post-bloom.

BUSHBERRIES, STRAWBERRIES, GRAPES: Apply 1 to 2 quarts/acre twice post-bloom.

MELONS, CUCUMBERS and SQUASH: Apply 1 to 2 quarts/acre 1 to 3 times after bloom.

MINT: Apply 1 quart/acre at 3 to 4 week intervals starting in seedling stage.

LETTUCE: Apply 1 quart/acre after thinning. Repeat with 2 quarts at 2 week intervals.

TOMATOES, PEPPERS, POTATOES, SWEET POTATOES and EGGPLANT: Apply 1 to 2 quarts/acre in 2 post-bloom applications.

ONIONS and GARLIC: Apply 1 to 2 quarts/acre at 1 to 4 week intervals throughout the growing season.

ORNAMENTALS and TURF: Apply 1 to 2 quarts/acre in 100 or more gallons water at monthly intervals.