

NOTE: This is a specimen label for electronic distribution. Refer to product label on container for specific directions for use.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS: CAUTION: Harmful if swallowed, inhaled, or contact is made with the skin or eyes. May cause severe irritation to eyes and moderate irritation to skin, lungs or stomach. Avoid contact with skin, eyes and clothing. Avoid breathing vapor or spray mist. Avoid using product in a manner as to directly, or through drift, expose workers or other persons. Always follow the precautionary statements on the label of the pesticide that is to be used with **CMR SILICONE SURFACTANT**.

ENVIRONMENTAL HAZARDS: Do not contaminate water when disposing of equipment washwaters.

STORAGE AND DISPOSAL

DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE OR DISPOSAL.

STORAGE: Keep container tightly closed. Do not allow water to be introduced to the contents of this container. Do not store near heat or open flame. Do not store or ship with oxidizers.

PRODUCT DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

GENERAL INFORMATION

CMR SILICONE SURFACTANT is a new generation surfactant based on organomodified siloxane technology for use in water-based pesticide formulations. **CMR SILICONE SURFACTANT** is a highly effective nonionic wetter/spreader/penetrant surfactant. Careful observation of spray application is recommended to determine proper and efficient pesticide application. This will optimize the pesticide and surfactant load levels.

While **CMR SILICONE SURFACTANT** has been proven to be a highly efficient surfactant, timing, weather conditions, methods of application, crop conditions, and/or mixture with other chemicals not specifically recommended are beyond the control of the seller. Therefore, the user should carefully observe spray deposits, rate efficiencies, compatibilities and effectiveness in initial application and adjust the adjuvant rates accordingly.

DIRECTIONS FOR USE

Pesticide manufacturers may recommend a rate of nonionic surfactant to be used with their product. Follow the pesticide label recommendations on use of surfactant. However, if the pesticide label does not have a specific recommendation for surfactant use, but does not prohibit the use of a nonionic surfactant, **CMR SILICONE SURFACTANT** should be applied at the rates listed below. Careful observation of spray performance in a trial application with the combination of chemicals to be applied is recommended to determine proper and efficient surfactant rates.

MIXING INSTRUCTIONS

(1) Fill tank to 2/3 to 3/4 full of water. If foaming is anticipated, add an anti-foaming agent before pesticide and **CMR SILICONE SURFACTANT**. (2) Add pesticide as directed by pesticide label. (3) Continue agitation. (4) Add **CMR SILICONE SURFACTANT** and continue filling tank while maintaining agitation.

FOR AGRICULTURAL SPRAYS



PRINCIPAL FUNCTIONING AGENTS:

A blend of polyether-polymethylsiloxane-copolymer and nonionic surfactant 100%

Calif. Reg. No. 1050775-50025-AA

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Immediately wash eyes with large amounts of water for at least 15 minutes, lifting the lower and upper lids. Seek medical attention immediately.

IF ON SKIN: Remove contaminated clothing and immediately wash contaminated area with large amounts of water for at least 15 minutes. If contact is made with the spray solution containing pesticides, follow the "Statement of Practical Treatment" on pesticide label. Seek medical attention.

IF SWALLOWED: If person is conscious, immediately give large quantities of water and induce vomiting, unless otherwise directed by label on pesticide(s). Do not induce vomiting or give fluids by mouth to an unconscious person. Seek medical attention immediately.

IF INHALED: Move person to fresh air and check to ensure adequate respiration. Seek medical attention immediately.

NET CONTENTS: 1 QUART

Manufactured by



CREATIVE MARKETING & RESEARCH, INC.
P.O. Box 35000 • Fresno, CA 93745-5000 • (559) 499-2100

* Registered TM of CREATIVE MARKETING & RESEARCH, INC.

AS A SPRAY ADJUVANT

With herbicides, defoliants, micronutrients: Apply at a rate of 6-16 ounces per 100 gallons of spray solution.

With insecticides, miticides, fungicides: Apply at a rate of 6-12 ounces per 100 gallons of spray solution.

For use with aerial applications: Apply at a rate of 12-16 ounces per 100 gallons of spray.

FOR USE WITH LIQUID FERTILIZATION

Apply **CMR SILICONE SURFACTANT** to liquid fertilizer/water solutions at the rate of 6-13 ounces per 100 gallons of mix. Not a plant food ingredient.

AS A SOIL WETTING AGENT

Golf course tees and greens: Apply at a concentration of 0.1% v/v or use the spray rate chart below.

<u>Spray rate/1000 sq. ft.</u>	<u>Amount of CMR SILICONE SURFACTANT in mix</u>
1.0 gal.	1/8 ounce
1.5 gal.	1/5 ounce
2.0 gal.	1/4 ounce

Repeat as needed to overcome hydrophobic soil condition. Irrigation is not needed, but will help incorporate the treatment into the soil.

For large turf areas such as golf fairways:

Boom sprayers: Apply at concentration of 0.05 - 0.1% or approximately 6-13 ounces per 100 gallons of tank mix/water.

Injection through irrigation: Adjust proportioning valve to inject at the ratio 1:1000, based on weekly application. To accommodate 100:1 proportioners, mix a 10% solution of product and water and inject at 100:1.

As a mulch, peat or potting soil wetting agent: Apply at a concentration of 0.05-0.1% v/v of water (6-13 ounces per 100 gallons). Repeat as needed.

AS A DEW CONTROL AGENT

CMR SILICONE SURFACTANT will prevent dew formation on turf areas such as tees and greens. Apply with a boom sprayer or hand sprayer. Use a concentration of 0.1% in water or approximately 13 ounces per 100 gallons every 7-10 days.

CONDITIONS OF SALE AND WARRANTY

Creative Marketing & Research, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated when used in accordance with the use directions under normal conditions. Creative Marketing & Research, Inc. neither makes, nor authorizes any agent or representative to make, any other warranties, express or implied, including fitness or merchantability.

The directions for use of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as timing and method of application, weather and crop conditions, presence of other materials, or other influencing factors, all of which are beyond the control of Creative Marketing & Research, Inc. and Seller. Buyer and user acknowledge and assume all risks and liability resulting from the handling, storage and use of this material not in strict accordance with directions given herewith. In no case shall Creative Marketing & Research, Inc. or the Seller be liable for consequential, special, indirect, or incidental damages or losses resulting from the handling or use of this product. The foregoing is a condition of sale by Creative Marketing & Research, Inc. and is accepted as such by the Buyer.