

# Crop-Set™

Apply Crop-Set at 8 oz/acre (240 mls/acre) or 600 mls/Ha. Dilute in a minimum of 10 gallons of water per acre (40 liters/acre) or 100 liters/Ha.

## CROP

Ganola (oilseed rape)  
Citrus  
Flower, leaf and stem veg.  
Fruit tree (less, citrus)

Melons, cucurbit  
Onion and other bulb crops  
Peas and Beans  
Potatoes (commercial)  
Potatoes (seed)  
Root veg. (carrots, etc.)  
Tomatoes, soft fruit

## TIME OF APPLICATION

Apply at rosette stage just prior to elongation of the stem.  
Apply between petal drop and 1 cm fruit; second application when fruit is golf-ball size.  
Apply at 5-6 leaf stage.  
Apply at 1-1.5 inches diameter fruit size and second when the fruit is about 1/2 the size of mature fruit.  
Apply soon after first flower.  
Apply at the 5-6 leaf stage.  
Apply at the 5-6 trifoliolate stage.  
Apply when largest tubers are 1-1.5 inches in diameter.  
Apply when stonions are swollen and hooked ends are formed.  
Apply at 5-6 leaf stage, plus 4-6 weeks after first application if in a stress situation.  
Apply at 10-30% of bloom. For multi-pick varieties, also apply after each picking.

## SHAKE WELL BEFORE USING

Net Weight: 1 gallon (9.5 Lb.) (4.31 kg) (3.78 Liter)

**STORAGE & PRECAUTIONS:** Keep container tightly sealed when not in use. Store in cool, dry place between 34° and 100°F (1° and 38°C). Do not store in direct sunlight. Do not store diluted product. Keep out of reach of children. Liquid should not be consumed. When used in conjunction with fungicides and insecticides the following precautions apply: Do not inhale mist. Wash with soap and water after use or skin contact. Prevent contamination of food, utensils and drinking water. Follow all governmental procedures for the disposal of the container; contact agricultural commissioner for disposal information.

*A source of micronutrients for improving plant growth and harvest yields in fruit, tuber, vegetable and seed production.*

## GUARANTEED ANALYSIS

|   |       |
|---|-------|
| Sulfur (S).....   | 1.50% |
| 1.50% Combined Sulfur (S)   |       |
| Copper (Cu).....  | 0.20% |
| 0.20% Water Soluble Copper (Cu)                                   |       |
| Iron (Fe).....  | 0.60% |
| 0.60% Water Soluble Iron (Fe)                                     |       |
| Manganese (Mn).....   | 1.50% |
| 1.50% Water Soluble Manganese (Mn)                                |       |
| Derived from: Manganese Sulfate, Copper Sulfate, Ferrous Sulfate. |       |

## ALSO CONTAINS NONPLANT FOOD INGREDIENTS

75.0% Bacterial fermentation product  
4.0% Plant Extract  
5.0% Yeast Extract

**Batch Number:**

**Expiration Date:**

Seller is not responsible for damage, loss or injury of any kind including, without limiting, lost profits. This product must be used as instructed. Express and implied warranties are disclaimed and sellers' responsibility shall not exceed the purchase price of the product.



# IMPROCROP LTD.

3031 Catnip Hill Pike • Nicholasville, KY 40356 • U.S.A. • Tel: (606) 887-3241 • Fax: (606) 887-3256

REV 7/02  
BT0018  
OREGON