

**NOTE: This is a specimen label for electronic distribution. Always refer to product label on container for specific directions for use.**

# FOLAR® 11-8-5

## GUARANTEED ANALYSIS 11-8-5

Total Nitrogen (N) .....	11.00%
3.35% Ammoniacal Nitrogen	
1.40% Nitrate Nitrogen	
6.25% Urea Nitrogen	
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> ) .....	8.00%
Soluble Potash (K <sub>2</sub> O) .....	5.00%
Copper (Cu) .....	0.05%
0.05% Chelated copper	
Iron (Fe) .....	0.10%
0.10% Chelated iron	
Manganese (Mn) .....	0.05%
0.05% Chelated manganese	
Zinc (Zn) .....	0.05%
0.05% Chelated zinc	

Derived from ammonium nitrate, ammonium phosphate, potassium nitrate, urea, copper ethylenediaminetetraacetate (EDTA), iron EDTA, manganese EDTA, and zinc EDTA.

WARNING: This product contains chemicals known to the State of California to cause cancer and birth defects or other reproductive harm.

## KEEP OUT OF REACH OF CHILDREN CAUTION

CAUTION: Causes eye irritation. In case of contact with the eyes, immediately flush with water for at least 15 minutes. If irritation persists, get medical attention. Do not swallow. Do not repack into unlabeled containers, or those normally used for food or drink.

FOLAR 11-8-5 provides nutrients for quick-response foliar feeding. It provides micronutrients that may be deficient in the soil or plant. It is compatible with organic pesticides to permit combination spraying.

**READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS.**

**If frozen during storage or shipment, heat to at least 70°F and agitate to redissolve crystals.**

## STATEMENT OF WARRANTY

NOTICE: Seller's guarantee shall be limited to the terms of the label. Subject thereto, the buyer assumes any risk to persons or property arising from the use or handling and accepts the product on these conditions.

**NET CONTENTS: 2.5 GALLONS / 9.5 LITERS**

NET WEIGHT: 10.25 lbs./gal. at 68°F

Manufactured by:



**CMR**

**CREATIVE MARKETING & RESEARCH, INC.**  
P.O. Box 35000 • Fresno, CA 93745-5000  
(559) 499-2100

\* Registered TM of CREATIVE MARKETING & RESEARCH, INC.

0185/0501(04)

# FOLAR 11-8-5

## DIRECTIONS FOR USE

For row or field crops, mix **FOLAR 11-8-5** in at least 20 gallons water per acre for ground sprayers or 10 gallons for aircraft. Apply to fruit trees or ornamentals in at least 100 gallons water per acre with conventional sprayers, 45-100 gallons with concentrate sprayers, or 10 gallons for aircraft. On waxy foliage or fruit, add sufficient spreader for smooth wetting to avoid spotting.

Use quantities indicated in the table below or follow recommendations of local agricultural authorities. Add spreader as needed for smooth wetting.

**USE PRECAUTIONS:** The high nitrogen (from 3 different sources) analysis is designed to stimulate early-season plant growth. Make test applications on a few plants if using combinations not known to be safe. Always add sufficient spreader to insure smooth wetting. Large droplets will run off young waxy foliage. Large droplets may also cause spotting and/or burning of leaf margins. Use only in equipment free of herbicide contamination. To avoid phytotoxicity, do not apply with copper or lime-sulfur fungicides, or during bloom.

**APPLES, APRICOTS, PEARS, NECTARINES, PEACHES, PLUMS, PRUNES, CHERRIES, ALMONDS, PECANS, WALNUTS, OLIVES, FIGS and AVOCADOS:** Apply 2 quarts/acre at first flush of spring growth. Repeat at post-bloom or as needed during the growing season.

**ALFALFA, CLOVER:** Apply 1 quart/acre as first spring growth starts and repeat after each cutting.

**BEANS, BARLEY, CORN, RICE, MILO, WHEAT, OATS, SAFFLOWER and SUGAR BEETS:** Apply 1 to 2 quarts/acre in early seedling stage and repeat at 2 to 3 week intervals.

**CITRUS:** Apply 2 quarts/acre at 2 to 3 week intervals as needed.

**COTTON:** Apply 2 quarts/acre after thinning. Repeat with 2 quarts at 2 week intervals until bloom.

**ARTICHOKES, CELERY, CARROTS, COLE CROPS, and SPINACH:** Apply 2 quarts/acre at 2 to 3 week intervals as needed.

**BUSHBERRIES, STRAWBERRIES and GRAPES:** Apply 2 quarts/acre at first foliage growth. Repeat at 2 to 3 week intervals until bloom.

**LETTUCE:** Apply 1 quart/acre after thinning. Repeat with 2 quarts at 2 week intervals.

**MELONS, CUCUMBERS and SQUASH:** Apply 1 quart/acre on first true leaves. Repeat at 2 week intervals with 2 quarts/acre until bloom.

**MINT:** Apply 1 quart/acre at 3 to 4 week intervals starting in seedling stage.

**TOMATOES, PEPPERS, POTATOES, SWEET POTATOES and EGGPLANT:** Apply 1 quart/acre on first true foliage. Repeat at 2 week intervals with 2 quarts/acre until bloom.

**ONIONS and GARLIC:** Apply 1 to 2 quarts/acre at 1 to 4 week intervals throughout the growing season.

**ORNAMENTALS and TURF:** Apply 1 to 2 quarts/acre in 100 or more gallons water at monthly intervals.