

SUPPLEMENTAL LABELING



Drexel

MSMA 6 Plus

MSMA Liquid Plus Surfactant

(EPA Reg. No. 19713-42)

FOR USE IN COTTON

**READ THE ENTIRE DREXEL MSMA 6 PLUS LABEL BEFORE PROCEEDING
WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABEL**

KEEP OUT OF REACH OF CHILDREN

CAUTION

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. This labeling must be in the possession of the user at the time of pesticide application.

COTTON

This product is useful for the control of weeds listed under *GENERAL INFORMATION* section and many similar weeds.

Application can be made: 1) Preplant or postplant up to cracking of soil before Cotton emergence using ground or aircraft equipment. 2) Postemergent, over the top, when Cotton is 3 to 6 inches high or up to early first square stage, whichever occurs first using ground or aircraft equipment. 3) Postemergent as a directed spray with ground equipment when Cotton is 3 inches high to first bloom.

Slight burning and a reddish discoloration of the Cotton leaf may occasionally occur following application, but the Cotton plant will develop normally.

DO NOT allow spray or spray drift to contact adjacent crops or injury will result. Apply only on still days when weather conditions DO NOT favor drift from areas being treated. Aircraft applications of this product should only be made by applicators experienced in use of herbicides, and application should be made in accordance with State and Federal regulations.

Note: Applications to Cotton in Florida should be confined to band treatments.

Pre-Plant Application or Post-plant up to Cracking Application: If planting of Cotton has been delayed and weeds have emerged, a single ground or aircraft application of this product can be made to prepared Cotton seedbeds or as a postplant treatment, but no later than initial cracking of soil. Cotton may be planted immediately following the pre-plant application. Mix at the rate of 2.66 pints of this product in 40 gallons of water for ground equipment or in 5 to 10 gallons of water for aircraft application per acre of Cotton.

Post-Emergent Applications Using Ground or Aircraft Equipment as an Over-the-Top Broadcast Spray when Cotton is 3 inches high until early first square stage as a salvage operation. **One application:** Use 2.5 pints in 40 gallons of water for ground application, or in 5 to 10 gallons of water for aircraft application per acre. **DO NOT** repeat this treatment. **Two applications:** Mix at the rate of 1 to 1.25 pints of this product in 40 gallons of water for ground equipment or in 5 to 10 gallons of water for aircraft application per acre of Cotton. If needed, a second or repeat application should be made 1 to 3 weeks after the first application. Apply only as a salvage operation. Apply only to healthy, rapidly growing Cotton, 3 inches high, but no later than 6 inches high or early square, whichever occurs first. Preference should be given to directed sprays. In order to minimize injury, the second application should be made as a directed spray when possible. **DO NOT** make more than two applications total of either DSMA or MSMA (or a combination) per season. Do not feed foliage to livestock or graze treated areas.

Postemergent Directed Spray Applications: Mix this product at the rate of 2.66 pints in 40 gallons of water per acre. For band applications, apply 1 gallon in 40 gallons of water per acre. For band applications, apply 1 gallon of above diluted spray per acre for each 1 inch band width to be treated of Cotton grown on 40 inch row spacing. Direct the spray solution to the base of the Cotton plant. Nozzles should be placed so as to avoid spraying the Cotton foliage, but to give good coverage of the weeds. If regrowth occurs, make a second application about 1 to 3 weeks after the first.

Make directed applications only when Cotton is 3 inches high to first bloom. Do not apply after first bloom. Do not feed foliage to livestock or graze treated areas.

All applicable directions, restrictions and use precautions on Drexel MSMA 6 PLUS label(s) are to be followed. This labeling contains supplemental instructions and restrictions for use of this product which may not appear on the Drexel MSMA 6 PLUS label. Follow the instructions carefully.

Manufactured By:



Drexel Chemical Company

P.O. BOX 13327, MEMPHIS, TN 38113-0327

SINCE 1972

42SUPP-0504
MSMA 6 PLUS