

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. May cause irritation of eyes, nose, throat and skin. Avoid contact with eyes, skin and clothing. Avoid breathing spray mist. Wash thoroughly after handling. Remove contaminated clothing and wash thoroughly before reuse. Do not contaminate feed and foodstuffs.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water. If so advised by a physician or Poison Control Center, induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF IN EYES: Immediately flush eyes with plenty of water and get medical attention.

IF ON SKIN: Immediately wash skin with plenty of soap and water while removing contaminated clothing and shoes. If irritation persists, seek medical attention.

IF INHALED: Remove victim to fresh air.


ENVIRONMENTAL HAZARDS

Do not contaminate water when disposing of equipment washwaters.

STATEMENT OF WARRANTY

NOTICE: Seller's guarantee shall be limited to the terms of the label. Subject thereto, the buyer assumes any risk to persons or property arising from use or handling and accepts the product on these conditions.

FOR AGRICULTURAL SPRAYS



BIO-DEGRADABLE
ACTIVATOR • BUFFERING AGENT • SPREADER

Calif. Reg. No. 1050775-50011-AA

KEEP OUT OF REACH OF CHILDREN

CAUTION

NET CONTENTS: 1 GALLON

Manufactured by:



CWR

CREATIVE MARKETING & RESEARCH, INC.

P.O. Box 2646
Fresno, CA 93745-2646

* Registered TM of CREATIVE MARKETING & RESEARCH, INC.

NO FOAM NUTRIENT FOLIAR SOLUTION 8-8-2 is a nutrient solution combined with a complete anionic/nonionic surfactant blend, **NO FOAM NUTRIENT FOLIAR SOLUTION 8-8-2** supplies major and minor nutrients and a complete surfactant system with emulsion stabilizer. It increases the efficiency of various agricultural chemicals. It insures quick wetting and uniform coverage and aids the performance of translocated and systemic agricultural chemicals. In areas where the water is alkaline, this product acidifies and buffers, tending to reduce hydrolysis of and increase the life expectancy of pesticides such as organic phosphates. **NO FOAM NUTRIENT FOLIAR SOLUTION 8-8-2** is designed to increase the absorption and utilization of micronutrients and is formulated from raw materials which will not tend to burn foliage.

GUARANTEED ANALYSIS

8 - 8 - 2	
Total Nitrogen (N)	8.00%
8% Ammoniacal Nitrogen	8.00%
Soluble Phosphoric Acid (P ₂ O ₅)	2.00%
Iron (Fe)	0.10%
Manganese (Mn)	0.05%
Zinc (Zn)	1.00%
Nitrogen and Phosphate from mono ammonium phosphate; Potash from potassium chloride; Iron, Manganese and Zinc from metallic sulfates.	
ADJUVANT	
PRINCIPAL FUNCTIONING AGENTS:	
Nonyl Phenoxy Polyethoxy Ethanol, Polydimethylsiloxane,	25.1%
Mono Ammonium Phosphate	74.9%
CONSTITUENTS INEFFECTIVE AS SPRAY ADJUVANT:	100.0%
TOTAL	100.0%

STORAGE AND DISPOSAL

DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE OR DISPOSAL.
STORAGE: Store in a dry location away from children, animals, foods, feeds and seeds. Handle in accordance with Precautionary Statements. In the event of spillage or leakage, soak up the material with absorbent clay, sand, sawdust or other absorbent material. Scrape up and dispose in accordance with Product Disposal.
PRODUCT DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
CONTAINER DISPOSAL: Triple rinse (or equivalent), then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration. If allowed by state and local authorities, dispose of containers by burning. If burned, stay out of smoke.

DO NOT STORE BELOW 32° FAHRENHEIT

DIRECTIONS FOR USE

APPLICATION DIRECTIONS: For row or field crops, apply in at least 20 gals. water per acre by ground and 40 gals. by aircraft. Apply to fruit trees in at least 100 gals. water/acre with conventional sprayers, 45-100 gals. with concentrate sprayers, and 10 gals. by aircraft. Agitate product in container first and during addition to water.

USE PRECAUTIONS: For best results, use this material in early stages of plant growth. Do not apply to fruit where scorching may occur or on crops not on this label. **NO FOAM NUTRIENT FOLIAR SOLUTION 8-8-2** is compatible with most crop protection chemicals. Do not use equipment contaminated with weed killers.

CAUTION: Tank mixtures of **NO FOAM NUTRIENT FOLIAR SOLUTION 8-8-2** with copper-based products can be phytotoxic due to increased solubility of copper. Test applications to part of the crop are advisable to assess potential phytotoxicity before full scale, commercial applications. The following crops are known to be sensitive to foliar applications of copper: broccoli, cabbage, celery, cucurbits, grapes, lettuce, nectarines, peaches, pears, plums, prunes, spinach, and strawberries.

PESTICIDE ADJUVANT FOR TERRESTRIAL USE SITES: Use 1 - 2 quarts product per 100 gallons of water to reduce the pH of alkaline waters used for pesticide applications.

FOLIAR NUTRITIONAL SPRAYS: Use the doses recommended below.

APPLES, APRICOTS, PEARS, PLUMS, PRUNES, NECTARINES, FIGS, PEACHES, CHERRIES, NUTCRACKS, & OLIVES: Apply 2 qts. per acre in delayed dormant and continue with several applications after blooming period. Use with **CAUTION** on smooth skinned fruits.

ALFALFA, BEANS, BARLEY, CORN, RICE & MILK: Apply 1 - 2 qts. per acre in seedling stage to early growth.

CITRUS: Apply 2 qts. per acre before or during spring flush of growth. Multiple applications are desirable. Apply 1 - 2 qts. per acre on non-bearing trees anytime before or during a flush of growth.

CELERY, CARROTS & COLE CROPS: Apply 1 - 2 qts. per acre 2 - 3 times in early growth.

COTTON: Apply 1 - 2 qts. per acre 1 - 3 times before full bloom.

BUSHBERRIES, STRAWBERRIES & GRAPES: Apply 1 - 2 qts. per acre during and after bloom.

LETTUCE: Apply 1 - 2 qts. per acre in 1 - 4 applications beginning at thinning time.

MELONS, CUCUMBERS & SQUASH: Apply 1 - 2 qts. per acre.

SUGARBEETS: Apply 1 - 2 qts. per acre in 1 - 2 applications from 6-leaf to 3 months old.

TOMATOES, PINEAPPLES, PEPPERS, POTATOES, SWEET POTATOES & EGG-PLANTS: Apply 1 - 2 qts. per acre in 1 - 2 applications.

TURF & ORNAMENTALS: Apply 1 - 2 qts. per acre at monthly intervals.

SHAKE WELL BEFORE USING

03510995(04)