

# SUPPLEMENTAL LABELING



Drexel



## **Simazine 90DF**

Herbicide

(EPA Reg. No. 19713-252)

**FOR WEED CONTROL IN CERTAIN CROPS AND ORNAMENTAL PLANTINGS.  
READ THE ENTIRE DREXEL SIMAZINE 90DF LABEL BEFORE PROCEEDING  
WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABEL**

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

**DIRECTIONS FOR USE**

**It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. This labeling must be in the possession of the user at the time of pesticide application.**

#### **TANK MIXTURE WITH ROUNDUP®**

This tank mixture is effective in Grape vineyards and in the following bearing or non-bearing tree crops for control of existing vegetation and for residual control of the annual Broadleaf and Grass weeds claimed for this product applied alone. This combination is also effective for partial control of Perennial weeds contacted by the spray mixture during application.

<b>Use this tank mix on these crops:</b>		
Almonds** (CA Only)	Grapefruit*** (CA, FL & TX)	Peaches**** <sup>2</sup>
Apples*	Grapes*	Pears*
Avocados* (CA & FL Only)	Lemons* (AZ & CA Only)	Pecans**
Cherries (Sour & Sweet****)	Macadamia nuts**	Plums**** <sup>1</sup>
Filberts**	Oranges*** (AZ, CA, FL & TX)	Walnuts**

**SPECIFIC DIRECTIONS:** Use the appropriate rate given on the Section 3 label for this product applied alone to the crop being treated. Add to the spray tank 1 to 5 qts. of Roundup per acre depending on weeds present and their growth stage. Also add an agriculturally approved non-ionic surfactant at 0.5% by volume of spray solution. Apply the mixture in 10 to 40 gals. of water per acre as a post-emergence spray to the weeds at the appropriate weed growth stage given on the Roundup label. Add this product to the spray tank first, then add Roundup. Provide constant agitation during mixing and application to keep the mixture in suspension. Refer to "APPLICATION PROCEDURES" section on the Section 3 label for further directions. Refer to labels of both herbicides for further directions, specific weeds controlled, precautions and limitations on each crop.

**USE PRECAUTIONS AND RESTRICTIONS (To avoid crop injury):** Take extreme care to avoid contact of herbicide solution, spray, drift or mist with foliage or green bark of trunk, branches, suckers, fruit, or other parts of trees or vines. Observe use precautions on both this product and Roundup labels for each crop involved.

\*Allow a minimum of 14 days between last application and harvest.  
\*\*Allow a minimum of 21 days between last application and harvest of these crops.  
\*\*\*Allow a minimum of 1 day between last application and harvest of these crops.  
\*\*\*\*Allow a minimum of 17 days between last application and harvest of these crops.  
<sup>1</sup>Plums and Sweet cherries: Limited to MO and states East of the Mississippi River, except TN.  
<sup>2</sup>Limited to AR, CA, LA, MO, OK, TX and states East of the Mississippi River. For CA, see specific directions in the section "ALMONDS, NECTARINES AND PEACHES (CA Only)"

**All applicable directions, restrictions and use precautions on Drexel SIMAZINE 90DF label(s) are to be followed. This labeling contains supplemental instructions and restrictions for use of this product which may not appear on the Drexel SIMAZINE 90DF label. Follow the instructions carefully.**



Manufactured By:

**Drexel Chemical Company**

P.O. BOX 13327, MEMPHIS, TN 38113-0327

**SINCE 1972**