

# SLUG-FEST

## ALL WEATHER FORMULA



*Lasts longer under wet conditions*

# ORCAL, Inc.

OR-CAL, Inc.  
29454 Meadowview Road  
Junction City, Oregon 97448

EPA Registration No. 71096-4  
EPA Establishment No. 52251-OR-005  
Patent No. 2,957,803

Molluscicide for Agricultural  
& Ornamental Usage

### 25% Metaldelhyde

**ACTIVE INGREDIENT:** Metaldelhyde (2,4,6,8-tetramethy-1,3,5,7-tetraoxycyclo-octane)..... 25%  
**INERT INGREDIENTS**..... 75%  
**TOTAL**..... 100%

*Contains 2.1 pounds Metaldelhyde per gallon.*

ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE LABEL ARE TO BE FOLLOWED. SEE DIRECTIONS FOR USE IN BOOKLET.

**KEEP OUT OF REACH OF CHILDREN  
WARNING AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

**SHAKE WELL**

**NET CONTENTS**

- 1 QUART
- 1 GALLON

### STATEMENT OF PRACTICAL TREATMENT

**If Swallowed:** Contact physician, Poison Control Center, or emergency clinic before inducing vomiting. Give water to drink and induce vomiting by touching back of throat with finger, or by administering syrup of ipecac. Do not induce vomiting or give anything by mouth to an unconscious person. If medical advice cannot be obtained, take person and product container to the nearest emergency treatment facility.  
**If Inhaled:** Remove person from exposure area.  
**If On Skin:** Wash skin with plenty of soap and water.  
**If In Eyes:** Wash eyes immediately with fresh water for at least 15 minutes. See a doctor if irritation persists.

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**WARNING.** May be fatal if swallowed. Avoid breathing spray mist. Do not get in eyes, on skin or on clothing. Wash thoroughly after use. Keep out of reach of children or domestic animals.  
**IMPORTANT: KEEP CHILDREN AND PETS OUT OF TREATED AREA UNTIL SPRAY HAS DRIED. THIS PESTICIDE MAY BE FATAL TO DOGS OR OTHER PETS IF EATEN.**

#### PERSONAL PROTECTIVE EQUIPMENT (PPE)

*Applicators and handlers must wear:* Coveralls over long-sleeved shirt and long pants, chemical-resistant or waterprool gloves, chemical resistant footwear plus socks, protective eyewear, chemical resistant apron when cleaning equipment, milking, or loading. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### USER SAFETY RECOMMENDATIONS

- Users should:**
- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
  - Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
  - Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## ENVIRONMENTAL HAZARDS

This product is toxic to birds and wildlife. Birds feeding on treated areas may be killed. For terrestrial uses, do not apply directly to water or areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water.

### STORAGE AND DISPOSAL

**Prohibitions:** Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

**Storage:**

Keep pesticides in original container in a locked place out of reach of children. Protect from freezing.

**Pesticide Disposal:**

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Disposal:**

Do not reuse container. Dispose of container in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### USE PRECAUTIONS

Read entire label before using. Apply directly to the soil area beneath plants as listed on this label. Do not allow spray to come in contact with parts of plant that may be used for food or feed purposes. The grower is responsible for residues on his crops, as well as for damages caused by drift from his property to that of others. Do not tank mix this product with other chemicals. Do not allow drift to areas occupied by unprotected humans or beneficial animals or onto adjoining food, fiber or pasture crops. *This product may be injurious to Daylilies and Clematis. Do not use on these plants.*

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**REENTRY:** Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

**EARLY ENTRY PPE:** PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: Coveralls over long-sleeved shirt and long pants; chemical resistant or waterproof gloves; chemical resistant footwear plus socks, and protective eyewear.

#### MIXING: Shake SLUG-FEST Colloidal 25 well before mixing.

Thoroughly mix spray solution before application. Use entire mixed solution at one time. Do not store mixed solution. Coagulation or mixture on sprayer screens can occur. Removal of screens from spray systems that operate at medium to high pressure is recommended.

**LAWNS AND ORNAMENTALS:** Such as Geraniums, Salvia, Primroses, Marigolds, Petunias, Pansies, Dahlias, Iris, Daisies, Begonias, Coleus, Ajuga and Dichandra.

**HOSE-ON TYPE APPLICATOR:** Thoroughly water areas to be treated. Use one ounce of SLUG-FEST Colloidal 25 for each two gallons of solution delivered by your applicator to cover approximately 200 sq. ft.

**SPRINKLING CAN:** Thoroughly water areas to be treated. Mix one ounce of SLUG-FEST Colloidal 25 to each two gallons of water and sprinkle over areas to be treated.

**WHERE SNAILS ARE A PROBLEM:** We suggest double rate of application on ornamentals only. **BEST RESULTS:** Apply SLUG-FEST Colloidal 25 in the cool of the evening.

## **TERRESTRIAL FOOD CROPS**

Alfalfa, Almond, Anise, Artichoke, Asparagus, Barley, Beans, Beets, Blackeyed Peas, Broccoli, Brussels Sprouts, Cabbage, Cantaloupes, Caraway, Carrots, Catinip, Cauliflower, Celery, Chikory, Chive, Clover, Collards, Comander, Corn, Cowpeas, Cucumbers, Currant, Dill, Eggplants, Endive, Garlic, Ginseng, Horse-radish, Kale, Kohlrabi, Leeks, Lentils, Lettuce, Melons, Muskmelons, Mustard, Mustard Greens, Oats, Okra, Onions, Oregano, Parsley, Parsnips, Peas, Peppermint, Peppers, Pimientos, Potatoes, Pumpkins, Radishes, Rhubarb, Rutabagas, Sage, Salsify, Savory, Shallots, Spear-mint, Spinach, Squash, Sugar Beets, Sweet Potatoes, Swiss Chard, Tomatoes, Turnips, Watermelon, Wheat, Seed Crops such as alfalfa, beans, clover, corn(sweet), grasses, peas, potato.

**SLUGS AND SNAILS:** Apply 2 to 4 quarts per treated acre in sufficient water for thorough coverage OR 5 to 10 oz. in ten gallons of water per treated 1,000 sq. ft. OR 1 to 2 oz. in two gallons of water per 200 sq. ft. Where snails are a problem, we suggest the higher rate of application. Thoroughly water areas to be treated before application. Apply only to soil surface around plants and in rows between plants. Use equipment that applies directly to the soil surface. Do not apply in such a manner as to cause drift of product onto plants. Do not apply to edible parts of plants. For best results, apply in the evening. Repeat as needed for control.

## **TREE CROPS**

Apples, Apricots, Avocados, Bananas, Cherries, Citrus fruits, Grapefruits, Lemons, Limes, Nectarines, Oranges, Peaches, Pears, Pecans, Plums, Stone fruits, Walnuts.

**SLUGS AND SNAILS:** Apply 2 to 4 quarts per treated acre in sufficient water for thorough coverage OR 5 to 10 oz. in ten gallons of water per treated 1,000 sq. ft. OR 1 to 2 oz. in two gallons of water per 200 sq. ft. Where snails are a problem, we suggest the higher rate of application. Make application when foliage and fruit are completely dry but soil is wet following an irrigation. Apply under trees, around irrigation pipes, farm buildings and any other place where slugs and snails may congregate. Use equipment that applies directly to soil surface. Do not apply in such a manner as to cause drift of product onto trees. Do not apply directly to trees or fruit. For best results, apply in the evening. Repeat as needed for control.

## **BERRY CROPS**

Blackberries, Blueberries, Boysenberries, Canberries, Dewberries, Gooseberries, Loganberries, Raspberries, Strawberries.

**SLUGS AND SNAILS:** Apply 2 to 4 quarts per treated acre in sufficient

water for thorough coverage OR 5 to 10 oz. in ten gallons of water per treated 1,000 sq. ft. OR 1 to 2 oz. in two gallons of water per 200 sq. ft. Where snails are a problem, we suggest the higher rate of application. Thoroughly water areas to be treated before application. Apply only to soil surface around plants and in rows between plants. Use equipment that applies directly to soil surface. Do not apply in such a manner as to cause drift of product onto plants. Do not apply to edible parts of plants. For best results, apply in the evening. Repeat as needed for control.

## **TERRESTRIAL NONFOOD CROPS**

### **SEED CROPS**

Beets, Cabbage, Carrots, Collards, Kale, Kohlrabi, Lettuce, Mustard, Radish, Spinach, Swiss Chard, Tobacco, Turnips.

**SLUGS AND SNAILS:** Apply 2 to 4 quarts per treated acre in sufficient water for thorough coverage.

Thoroughly water areas to be treated before application. Apply only to soil surface around plants and in rows between plants. Use equipment that applies directly to soil surface. Do not apply in such a manner as to cause drift of product onto plants. For best results, apply in the evening. Repeat as needed for control.

### **GREENHOUSE FOOD CROP**

Asparagus, Beans, Beets, Cabbage, Carrots, Celery, Corn, Cucumbers, Eggplant, Endive, Lettuce, Melons, Onions, Peas, Peppers, Potatoes, Radish, Spinach, Squash, Tomatoes, Turnips.

**SLUGS AND SNAILS:** Apply 2 to 4 quarts per treated acre in sufficient water for thorough coverage OR 5 to 10 oz. in ten gallons of water per treated 1,000 sq. ft. OR 1 to 2 oz. in two gallons of water per 200 sq. ft. Where snails are a problem, we suggest the higher rate of application. Water or irrigate areas to be treated before application. Apply only to soil surface around plants. In rows between plants and under benches in greenhouses and around plants, any place where slugs and snails may congregate. Use equipment that applies directly on soil surface. Do not apply in such a manner as to cause drift of product onto plants. Do not apply to plants or edible parts of plants. For best results, apply in the evening. Repeat as needed for control.

## **GREENHOUSE NONFOOD CROPS**

### **ORCHIDS, ORNAMENTAL PLANTS**

**SLUGS AND SNAILS:** Apply 2 to 4 quarts per treated acre in sufficient water for thorough coverage OR 5 to 10 oz. in ten gallons of water per treated 1,000 sq. ft. OR 1 to 2 oz. in two gallons of water per 200 sq. ft.

Where snails are a problem, we suggest the higher rate of application. Water or irrigate areas to be treated before application. Apply only to soil surface around plants, in rows between plants and under benches in greenhouses and around plants; any place where snails and slugs might congregate. Do not apply directly to plants unless prior experience has shown the plant to be tolerant to this product. Most plants are tolerant, however, some slight tip burning has been noticed on extremely tender orchids.

**DOMESTIC AND NONDOMESTIC OUTDOOR DOMESTIC DWELLINGS, FARM BUILDINGS, LATH HOUSES, ORNAMENTAL PLANTS.**

**SLUGS AND SNAILS:** Apply 2 to 4 quarts per treated acre in sufficient water for thorough coverage OR 5 to 10 oz. in ten gallons of water per treated 1,000 sq. ft. OR 1 to 2 ozs in 2 gallons of water per treated 200 sq. ft. Where snails are a problem we suggest the higher rate of application.

Thoroughly water areas to be treated before application. Use equipment that applies directly to the soil surface. Apply evenly over infested areas and along flower beds adjacent to border or ground cover plants. Also apply to ground at base of fences and walls in shady, protected locations and any other place where slugs and snails may congregate.

Do not apply in such a manner as to cause drift of product onto edible plants or plant parts. Do not apply directly to plants unless prior experience has shown the plant to be tolerant, however, some slight tip burning has been noticed on extremely tender orchids. For best results, apply in the evening. Repeat as needed for control.

**LAWNS (INCLUDING DICHOINDRA) ORNAMENTAL GRASSES, TURF, NEWLY SEEDED GRASS GROWN FOR SEED (prior to canopy closure)**

**SLUGS AND SNAILS:** Apply 2 to 4 quarts per treated acre in sufficient water for thorough coverage OR 5 to 10 oz. in ten gallons of water per treated 1,000 sq. ft. OR 1 to 2 oz. in two gallons of water per 200 sq. ft. Where snails are a problem, we suggest the higher rate of application. Uniformly broadcast directly on lawn or turf. Withhold heavy watering for at least 2 days after treatment. Keep children and pets off the treated area until spray has dried.

**CONDITIONS OF SALE — LIMITED WARRANTY AND LIMITATIONS OF LIABILITY AND REMEDIES**

The directions on this label are believed to be reliable and should be followed carefully. Insufficient control of pests and/or injury to the crop to which the product is applied may result from the occurrence of extraordinary or unusual weather conditions, the failure to follow the label directions, or good application practices, all of which are beyond the control of OR-CAL, Inc. or seller. In addition, failure to follow label directions may cause injury to crops, animals, man or the environment. OR-CAL, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purpose referred to in the directions for use subject to the factors noted above which are beyond the control of OR-CAL, Inc. OR-CAL, Inc. makes no other warranties or representations of any kind, express or implied, concerning the product, including no implied warranty of merchantability or fitness for any particular purpose. The exclusive remedy against OR-CAL, Inc. for any cause of action relating to the handling or use of this product is a claim of damage and in no event shall damages or any other recovery of any kind against OR-CAL, Inc. exceed the price of the product which causes the alleged loss, damage, injury, or other claim. OR-CAL, Inc. shall not be liable and any and all claims against OR-CAL, Inc. are waived, for special, indirect, incidental, or consequential damages or expense, of any nature, including but not limited to, loss of profits, or income, whether or not based on OR-CAL, Inc. negligence, breach of warranty, strict liability in tort or any other cause of action. OR-CAL, Inc. and the seller offer this product and the buyer and user accept it, subject to the foregoing conditions of sale and limitation of warranty, liability and remedies.